

# ADVANCED ORTHOPAEDIC SPECIALISTS



Dr. Chris Arnold

Dr. David Yakin

Dr. Terry Sites

Dr. Mark Powell Dr. Kenton Hagan

# Postoperative Rehabilitation Following Elbow Arthroscopy (Posterior Compartment/Valgus Extension Overload)

# **Phase I: Immediate Motion Phase**

Goals: Improve/regain full range of motion
Decrease pain/inflammation
Retard muscular atrophy

# A. Day 1 to 4

- Range of motion to tolerance (Extension/flexion and supination/pronation) \*Often full elbow extension is not capable due to pain
- · Gentle overpressure into extension
- · Wrist flex/ext stretches
- Gripping exercises (putty)
- · Isometrics wrist ext/flex
- · Isometrics elbow ext/flex
- · Compression dressing, ice 4-5 times daily

## B. Day 5 to 10

- · Range of motion exercises to tolerance (at least 10-100 degrees)
- · Overpressure into extension
- · Joint mobilization to reestablish ROM
- · Wrist flex/ext stretches
- · Continue isometrics
- · Continue use of ice, compression to control swelling

## C. Day 11 to 14

- Range of motion exercises to tolerance (at least 20-90 degrees)
- · Overpressure into extension (3-4 times daily)
- · Continue joint mobilization techniques
- Initiate light dumbbell program (PRE's) Biceps, triceps, wrist flex/ext, sup/pronators

# **Phase II: Intermediate Phase**

Goals: Improve strength/power/endurance

Increase range of motion Initiate functional activities

# A. Week 2 to 4

- · Full range of motion exercises (4-5 times daily)
- · Overpressure into elbow extension
- · Continue PRE program for elbow and wrist musculature
- Initiate shoulder program (esp ER, RTC)
- · Continue joint mobilization
- · Continue ice post-exercise

# B. Week 4 to 7

- · Continue all exercises listed above
- Initiate light upper body program
- · Continue use of ice post-activity

# **Phase III: Advanced Strengthening Program**

**Goals:** Improve strength/power/endurance
Gradual return to functional activities

## Criteria to enter Phase III

- 1. Full non-painful range of motion
- 2. Strength 75% > of contralateral side
- 3. No pain or tenderness

#### A. Week 8 to 12

- · Continue PRE program for elbow and wrist
- · Continue shoulder program
- · Continue stretching for elbow/shoulder
- · Initiate interval program and gradually return to sport activities